

Prepared by Alex Buchanan, previously of the Tasmanian Herbarium

Woodland dominated by *Eucalyptus amygdalina* and *E. viminalis*

The most widespread community on the dry sandy soils. In some areas, *E. amygdalina* is dominant, in others *E. viminalis* takes over, while elsewhere, the two species are co-dominant.

FERNS		<i>Comesperma volubile</i>	
<i>Pteridium esculentum</i>		<i>Conospermum hookeri</i>	e
MONOCOTYLEDONS		<i>Correa reflexa</i> var. <i>reflexa</i>	
<i>Burchardia umbellata</i>		<i>Dampiera stricta</i>	
<i>Deyeuxia quadriseta</i>		<i>Dillwynia glaberrima</i>	
<i>Hypolaena fastigiata</i>		<i>Dodonaea viscosa</i>	
<i>Lepidosperma concavum</i>		<i>Epacris impressa</i>	
<i>Lomandra longifolia</i>		<i>Erica lusitanica</i>	
<i>Patersonia fragilis</i>		i	
<i>Schoenus lepidosperma</i>		<i>Eucalyptus amygdalina</i>	e
<i>Xanthorrhoea australis</i>		<i>Eucalyptus sieberi</i>	e
<i>Xyris tasmanica</i>	e	<i>Eucalyptus viminalis</i> subsp. <i>viminalis</i>	
DICOTYLEDONS		<i>Euryomyrtus ramosissima</i>	
<i>Acacia genistifolia</i>		<i>Exocarpos cupressiformis</i>	
<i>Acacia longifolia</i> subsp. <i>sophorae</i>		<i>Gompholobium huegelii</i>	
<i>Acacia suaveolens</i>		<i>Hibbertia riparia</i>	
<i>Acacia terminalis</i>		<i>Leptospermum scoparium</i>	
<i>Acrotriche serrulata</i>		<i>Leucopogon collinus</i>	
<i>Allocasuarina littoralis</i>		<i>Leucopogon ericoides</i>	
<i>Allocasuarina monilifera</i>	e	<i>Monotoca elliptica</i>	
<i>Amperea xiphoclada</i>		<i>Monotoca scoparia</i>	
<i>Aotus ericoides</i>		<i>Olearia ramulosa</i>	
<i>Astroloma humifusum</i>		<i>Persoonia juniperina</i>	
<i>Banksia marginata</i>		<i>Philothea virgata</i>	
<i>Bossiaea cinerea</i>		<i>Phyllota diffusa</i>	e
<i>Brachyloma ciliatum</i>		<i>Pimelea linifolia</i>	
<i>Calytrix tetragona</i>		<i>Platylobium triangulare</i>	
<i>Cassinia aculeata</i>		<i>Pultenaea tenuifolia</i>	
<i>Cassytha glabella</i>		<i>Styphelia adscendens</i>	
<i>Cassytha melantha</i>		<i>Tetratheca labillardierei</i>	
<i>Cassytha pubescens</i>			

Open woodland of *Eucalyptus ovata* in damp hollows

Scattered low trees over a low shrub layer. This area is partially inundated during winter rainy periods, drying in summer.

FERNS		DICOTYLEDONS	
<i>Lindsaea linearis</i>		<i>Allocasuarina littoralis</i>	
<i>Selaginella uliginosa</i>		<i>Allocasuarina monilifera</i>	e
MONOCOTYLEDONS		<i>Aotus ericoides</i>	
<i>Baumea acuta</i>		<i>Dillwynia glaberrima</i>	
<i>Eriochilus cucullatus</i>		<i>Epacris impressa</i>	
<i>Hypolaena fastigiata</i>		<i>Eucalyptus ovata</i> var. <i>ovata</i>	
<i>Lepidosperma filiforme</i>		<i>Hakea teretifolia</i> subsp. <i>hirsuta</i>	
<i>Lepidosperma longitudinale</i>		<i>Leptospermum scoparium</i>	
<i>Leptocarpus tenax</i>		<i>Melaleuca gibbosa</i>	
<i>Lomandra longifolia</i>		<i>Persoonia juniperina</i>	
<i>Schoenus lepidosperma</i>		<i>Stylidium graminifolium</i>	

Open forest of *Eucalyptus sieberi*

Trees to about 20m tall over open shrub layer; restricted to a small area near the NW corner of the reserve.

MONOCOTYLEDONS

Danthonia sp.*Gahnia radula**Hypolaena fastigiata**Lepidosperma concavum**Lepidosperma longitudinale**Lomandra longifolia*

DICOTYLEDONS

*Acacia genistifolia**Acacia myrtifolia**Acacia suaveolens**Allocasuarina littoralis**Aotus ericoides**Epacris impressa**Eucalyptus amygdalina* e*Eucalyptus sieberi* e*Eucalyptus viminalis**Exocarpos cupressiformis**Leptospermum scoparium**Monotoca elliptica**Olearia ramulosa**Persoonia juniperina****Allocasuarina littoralis* forest**

Trees to about 20m with sparse understorey. The listed eucalypts are occasionally co-dominant.

FERNS

*Lindsaea linearis**Pteridium esculentum*

MONOCOTYLEDONS

*Austrostipa pubinodis**Burchardia umbellata**Gahnia radula**Hypolaena fastigiata**Lepidosperma concavum**Lepidosperma filiforme**Leptocarpus tenax**Lomandra longifolia*

DICOTYLEDONS

*Acacia myrtifolia**Acacia suaveolens**Allocasuarina littoralis**Banksia marginata**Epacris impressa**Eucalyptus amygdalina* e*Eucalyptus viminalis**Hibbertia riparia**Leptospermum scoparium**Leucopogon collinus**Monotoca elliptica***Wet areas**

On the margins of eucalypt-dominated communities. Saltmarsh species still to be listed completely.

MONOCOTYLEDONS

Lachnagrostis sp.*Apodasmia brownii**Baumea juncea**Centrolepis strigosa**Festuca arundinacea* i*Holcus lanatus* i*Ficinia nodosa**Juncus kraussii**Juncus pallidus**Poa poiformis*

DICOTYLEDONS

*Acacia dealbata**Acacia melanoxylon**Acacia verticillata* subsp. *verticillata**Actites magalocarpa**Brachyscome graminea**Conyza sumatrensis* i*Erica lusitanica* i*Goodenia ovata**Lawrenzia spicata**Leontodon saxatilis* i*Leptinella longipes**Lobelia anceps**Melaleuca ericifolia**Melaleuca squarrosa**Olearia lirata**Plantago coronopus* i*Pomaderris apetala* subsp. *apetala**Pultenaea dentata**Rubus* blackberry aggregate i*Vellereophyton dealbatum* i

The point at the end of the track, the old ford

MONOCOTYLEDONS

Aira elegantissima i
Festuca arundinacea i
Isolepis levynsiana ?i
Romulea rosea var. *australis* i
Vulpia bromoides i

DICOTYLEDONS

Acetosella vulgaris i
Atriplex sp.
Cerastium glomeratum i

Crassula spp.
Erodium cicutarium i
Geranium sp.
Moenchia erecta i
Oxalis pes-caprae i
Rumex brownii
Samolus repens
Trifolium cernuum i
Trifolium dubium i
Trifolium subterraneum i

Alex Buchanan, Dick Burns